CULTIVATION **RECOMMENDATION**

Agria.

Maincrop, yellow fleshed, processing organic potato variety with a good yield and a very stable quality.

- **Excellent frying quality**
- **Multiple virus resistances**
- **Good storability**

Organic

Processing
French fries
Crisps
Convenience

Retail
Fresh

Traditional



characteristics

General characteristics

Breeder: Böhm
Crossing: QUARTA x SEMLO
Breedersright: Expired
Seedling: BM 77-0945

Plant and tuber characteristics

Maturity: maincrop / 5
Light sprout colour: blue violet / BV
Flower colour: white / W
Number of berries: none / 8
Foliage - initial development: quite fast / 7
Foliage - final development: good / 8
Foliage - firmness: quite firm / 7
Skin colour: yellow / Y
Flesh colour: yellow / 7
Skin finish: average / 6
Tuber shape: oval long / OL
Regularity of tuber shape: regular / 7
Tuberisation: quite low / 5
Tuber size: large / 8
Regularity of size: regular / 7
Dormancy: long / 8

Sensitivity and quality

Secondary growth: quite sensitive / 6
Black spot (internal bruising): quite sensitive / 6
Growth cracks: quite sensitive / 6,5
Harvest damage: sensitive / 5
Metribuzin - pre-emergence: slightly sensitive / 7,5
Metribuzin - post-emergence: slightly sensitive / 7,5
Bentazon: quite sensitive / 6,5
Ethylene: 5% to 15% more tubers / 6
Underwater weight: 402
Dry matter content: 21,8%
Cooking type: floury / BC
Discolouration (after cooking): slightly sensitive / 7
Raw discolouration: slightly sensitive / 7
Fries quality: excellent / 8
Dry matter distribution: average / 6,5
Crisping quality: good / 7
TGA-level: 2,7

Agrico has taken all reasonable care in producing this variety description. However, please note that any information may be subject to change and Agrico accepts no responsibility for errors or omissions. In no event shall Agrico be liable for damages arising out of or related to the information contained within this description.

AGRICO

CULTIVATION RECOMMENDATION

How to grow

Fertilization [1]

Nitrogen: standard advice - 10%.
Phosphate: standard advice.
Potassium: standard advice + 10%.

Seed treatment

A short heat shock or proper pre-sprouting is recommended.Proper pre-sprouting after taking off the top sprout can give a positive result on the tuber set.

Planting distance

28/35 mm: 25 cm (54.000 plants/ha)
35/55 mm: 30 cm (45.000 plants/ha)
50/55 mm: 40 cm (33.000 plants/ha)

Weeds

After emerge, a standard use of metribuzin (Sencor) can give some foliage reaction, but rarely any yield reduction.

Fungicides

A preventive spraying schedule against late blight is advised.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least three weeks after haulm killing before harvest.

Storage - .

Preferred storage temperature is 5 ºC. For processing purposes, do not store below 7 ºC. Suitable for long term storage.



Say potato, say Agrico.

Resistances

Virus - Leaf roll: slightly susceptible / 6
Virus - Yn: slightly to not susceptible / 8
Virus - Yntn (tuber): slightly susceptible / 7
Spraing: slightly susceptible / 7
Late blight - foliage: susceptible / 5,5
Late blight - tuber: slightly susceptible / 7
Fusarium: slightly susceptible / 6,5
Common scab: very susceptible / 4,5

Powdery scab: very susceptible / 3
Black dot: - / -
Wart disease F1: slightly susceptible / 9
Wart disease F6: quite susceptible / 6
Wart disease F18: - / -
Golden nematode Ro 1/4: resistant / 9
Golden nematode Ro 2/3: - / -
White nematode Pa 2: susceptible / 2
White nematode Pa 3: susceptible / 2

Agrico • Duit 15 • 8305 BB Emmeloord • The Netherlands +31 (0)527 63 99 11 info@agrico.nl agricopotatoes.com